

## <u>Pre-Treatment Instructions for PRP Breast Lift</u>

Avoid the following for at least 1 week prior to your procedure:

- NSAIDS (Ibuprofen, Motrin, Aleve, Naprosyn, Aspirin)—Tylenol is allowed
- Certain nutritional supplements (Gingko, Garlic, Vitamin E, Vitamin A, Flax
   Oil, Curcumin, and other anti-inflammatory nutrients)
- Systemic Steroids (prednisone, dexamethasone) -Alcohol and Cigarettes

  Hydrate well the day before and the day of the procedure for ease of blood draw.

  We recommend using Arnica the day of and two days after the procedure to decrease the risk of bruising.

## On the Treatment Day:

Blood is drawn and PRP is processed

A topical anesthetic is applied to the treatment area for 20-30 minutes.

Fillers are injected when a defect is being augmented (indentations, inverted nipples).

PRP and PPP (Platelet Poor Plasma) are then injected into multiple areas of the

breasts to facilitate the rejuvenation and regeneration of tissue.

There may be redness, swelling, bruising, and a general "tight feeling" immediately post-procedure and for a few days after.

826 Main Street, Klamath Falls, OR, 97601 541-887-8229 Almondaesthetic.com



## Post-Treatment Instructions for PRP Breast Lift

- Expect mild inflammation, redness, and swelling for a few days.
- Bruising may last several weeks. You may apply arnica gel or cream to the area to help. You can continue to take Arnica, 5 pellets 3 times a day until bruising clears.
- For at least 1 week, avoid NSAIDS (Ibuprofen, Motrin, Aleve, Naprosyn,
   Aspirin), certain nutritional supplements (Gingko, Garlic, Vitamin E, Vitamin
   A, Flax Oil, Curcumin, and other anti-inflammatory nutrients), systemic
   steroids (prednisone, dexamethasone), alcohol and cigarettes.
- Eat healthy and hydrate very well (at least 64 ounces of water daily).
- A sports bra is recommended to wear for the first few days post-treatment for comfort reasons.
- You may want to wear 2 sports bras when exercising post-treatment for a few days for extra support.
- If PDO threads were used, avoid strenuous aerobic exercise for 1-2 weeks to allow healing time.

826 Main Street, Klamath Falls, OR, 97601 541–887–8229 Almondaesthetic.com



• If fillers were used, you will want to sleep upright and not put any pressure on this area of your breasts for 1-2 weeks.

826 Main Street, Klamath Falls, OR, 97601 541-887-8229 Almondaesthetic.com